

Zea mays yield prediction using satellite information and a simulation model

Di Bella,C.M.¹
Magrin,G.O.¹
Boullon,D.R.¹
Grondona,M.O.¹
Rebella,C.M.¹

¹Centro de Investigaciones de Recursos Naturales-Instituto de Clima y Agua-INTA Castelar
Los Reseros y Las Cabañas S/N (1712)- Castelar, Buenos Aires, Argentina
(e-mail: imagen@castelar.gov.ar)

Crop yield prediction represents an important instrument for making economical and technological decisions in agricultural production. It is also important for Governmental Organizations to analyze the annual market situation, specially those assigned to exportation.

Several studies demonstrated the advantages of satellite information and crop models to predict crop yield in extensive agricultural areas. The temporal resolution and low cost of satellite information derived from NOAA AVHRR images represents an advantage. However, it is difficult to distinguish land use with the spatial resolution obtained from these images. On the other hand, crop models, considering crop functioning have a high predictive capacity; but some input variables are not always available, so calibration works for local conditions are required.

The main objective of this study was to compare the predictive behavior of satellite information and a crop model (CERES-Maize) concerning the interannual variation of maize yields in the Pampas Region of Argentina.

We selected as study areas 5 counties of Buenos Aires Province with similar crop management conditions. In each county we processed the digital information derived from monthly NOAA AVHRR GAC images of South America and ran the Ceres-Maize model for the period 1981-1990.

For each study area we correlated the mean monthly NDVI (Normalized Difference Vegetation Index) from October to March with observed yields of maize. To run the model we considered a complete cycle hybrid, Dekalb 3S41, and the crop management that characterizes the whole region. Initial soil water content was set according to the occurred rainfall and

the water accumulated in the soil during 4 months before planting.

January NDVI values showed the better association with the observed yields for each county ($R^2=0.59$) with a mean error of 3.91 qq/ha (**Figure 1**). There was no interaction NDVI-county, but a significant effect of the county. In this way we obtained five curves, one for each county, with the same slope but different y-intercepts. The differences observed in the y-intercepts could be related to regional climatic variability.




Figure 1: Relationship between observed and estimated yields (qq/ha) by January NDVI derived from NOAA-AVHRR GAC images. (-) 1:1 •

CERES-Maize predicted yields with a $R^2=0.76$ (**Figure 2**) and a mean error of 3.29 qq/ha, which represents an error of 8.2 % related to the mean observed data (40.37 qq/ha).




Figure 2: Linear correlation between observed and estimated yields (qq/ha) using the CERES-Maize model.

Comparing the results obtained with both methods, we observed that crop model simulation explained better the interannual variation of observed maize yields ($r^2=0.76$ vs $r^2=0.59$) than monthly NDVI. Also, comparing the Mean Absolute Error (MAE) and the Root Mean Square Error (RMSE) of both methods, Ceres-Maize performance was better in estimating maize yields (Table 1).

	CERES-Maize	NOAA-AVHRR
Obs. Mean	40.36	40.36
Est. Mean	40.62	40.36
N	45	45
MAE	3.29	3.91
RMSE	4.24	5.94
r²	0.76	0.59

Table 1 : Comparative statistics parameters. Obs. Mean = Average of observed yields. Est. Mean = Average of estimated yields. N = Number of cases. MAE = Mean absolute error. RMSE = Root mean square error. r^2 = Regression coefficient.

The inputs necessary to run these models are not always available on a daily basis for the whole area. The satellite data, however, resumes in one monthly image the information needed to make the prediction.

We concluded that using satellite information together with other meteorological indexes (i.e. drought index) or as input for simulation models will improve maize yields prediction.

BIBLIOGRAPHY

Allen,DM (1974): The relationship between variable selection and data augmentation in a method of prediction. *Technometrics* 16, 125-129.
Box,EO; Holben,BN and Kalb,V (1989): Accuracy of the AVHRR vegetation index as a predictor of biomass, primary productivity and net CO₂ flux. *Vegetatio* 80: 71-89.
Delecolle,R and Guerif,M (1988): Introducing spectral data into a plant process model for improving its prediction ability. *Proceedings of*

the 4th International Colloquium on Spectral Signatures of Objects in Remote Sensing held at Aussois, France, pp. 18-22.

Estadísticas Junta Nacional de Granos. Período 1971-1990.

Hodges,T; Botner,D; Sakamoto,C and Hays Haug,J (1987): Using the Ceres-Maize model to estimate production for the U.S. Cornbelt. *Agricultural and Forest Meteorology* 40: 293-303.

Hayes, MJ and Decker, WL (1990): Utilising satellite data within a corn production assessment system for the United States Corn Belt.

Jones and Kiniry (1986): CERES Maize. A simulation model of maize growth and development. Texas A&M University Press. College Station.

Kennedy,P (1989): Monitoring the vegetation of Tunisian grazing lands using normalized difference vegetation index. *Ambio* 18: 119-123.

Kerdiles, H; Pujol,GC; Rebella,CM and Seguin,B (1995): Vegetation monitoring and yield prediction from NOAA-AVHRR data in the Pampas Region, Argentina. (Enviado for publication).

Liu,WTH; Botner,DM and Sakamoto,CM (1989): Application of Ceres-Maize model to yield prediction of a Brazilian maize hybrid. University of Missouri Agricultural Experiment Station Journal Series N° 10526. Elsevier Science Publishers B.V.,300-312.

Maas, SJ (1988): Using satellite data to improve model estimates of crop yield. *Agronomy Journal* 80, 655-662.

Magrin, GO (1994). Aplicaciones de Ceres-Wheat en la Región Pampeana Argentina. Actas de la II Reunión Nacional de Trigo y Cereales Menores. 17 al 19 de Agosto. Santa Cruz de la Sierra, Bolivia. (En prensa).

Magrin,GO; Hall, AJ and Travasso,MI (1995): An empirical approximation to rules for initializing Ceres-Wheat for use in yield prediction of county-seized reporting districts. Presented at: International Congress on modelling and simulation 1995. Newcastle, Australia, 27-30/11/95.

Magrin, GO (1985): Modèles de prévisions de rendiments. Bilan de quatre années de prévisions ITCF sur blé. *Perspectives Agricoles* n° 92.

Philipson,WR and Teng,WL (1988): Operational Interpretation of AVHRR vegetation indices for world crop information. *Photogrammetric Engineering and Remote Sensing* 54, pp. 55-59.

Prince,SD (1990): High temporal frequency remote sensing of primary production using NOAA AVHRR. In: Applications of remote sensing in agriculture butterworks (eds: Seven,MD & Clark,JA), London, pp. 169-183.

Ritchie,JT (1986): Ceres-Maize. A simulation model of maize growth and development. Chapter 1: The Ceres -Maize model. Texas A&M University Press, Texas, pp. 3-7.

Tucker,CJ; Vanpraet,CL; Sharman,MJ and Van Ittersum,G (1985): Satellite remote sensing of total herbaceous biomass production in the Senegalese Sahel: 1980-1984. *Remote Sensing of Environment* 17: 233-249.

Wiegand,CL; Richardson,AJ; Jackson,RD; Pinter Jr.,PJ; Asse,JK; Smika,DE; Lautenschlager,LF and McMurtry III,JE (1986): Development of Agrometeorological Crop Model inputs from remotely sensed information. *IEEE Transactions on Geoscience and Remote Sensing* 24, pp. 90-97.

Willmott, CJ (1981): Some comments on the evaluation of model performance. *Bulletin American Meteorological Society* 63, 1309-1313.